

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

THIS PAGE BLANK (USPTO)

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/68, 37/02, 47/48		A1	(11) International Publication Number: WO 94/27613
			(43) International Publication Date: 8 December 1994 (08.12.94)
(21) International Application Number: PCT/AU94/00274		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FL, GB, GE, HU, JP, KG, KP, KR, KZ, LK, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 24 May 1994 (24.05.94)		Published With international search report. With amended claims.	
(30) Priority Data: 08/064,873 24 May 1993 (24.05.93) US			
(71) Applicant (for all designated States except US): BIOTECH AUSTRALIA PTY. LIMITED [AU/AU]; 28 Barcoo Street, Roseville, NSW 2069 (AU).			
(72) Inventors; and (75) Inventors/Applicants (for US only): RUSSELL-JONES, Gregory, John [AU/AU]; 23 Greenfield Avenue, Middle Cove, NSW 2068 (AU). WESTWOOD, Steven, William [AU/AU]; 6/98 Alt Street, NSW 2131 (AU).			
(74) Agents: STEARNE, Peter, Andrew et al.; Davies & Collison, 1 Little Collins Street, Melbourne, VIC 3000 (AU).			
(54) Title: VITAMIN B₁₂ MEDIATED ORAL DELIVERY SYSTEMS FOR GCSF AND EPO			
(57) Abstract			
<p>The invention describes complexes between VB₁₂ analogues and either GCSF or EPO that retain both significant affinity for intrinsic factor (IF) in the VB₁₂ portion of the complex and significant bioactivity of the GCSF or EPO portion of the complex. The invention also concerns a process for the synthesis of these complexes. This is achieved at least in part, by using a spacer compound, which is linked covalently between the VB₁₂ portion and the GCSF or EPO. The complexes preferably have the formula V-X-A-Y-Z wherein V is vitamin B₁₂ or a vitamin B₁₂ analogue, or derivative, bonded to X either through a carboxylate group pendant to the corrin nucleus of VB₁₂ or through the central cobalt atom or to a function group introduced onto the VB₁₂ molecule, X is selected from: -NHNH-, -NH-, -O-, -S-, -SS- or -CH₂-, and A is an optionally substituted, saturated or unsaturated, branched or linear, C₁₋₅₀alkylene, cycloalkylene or aromatic group, optionally with one or more carbons within the linear chain being replaced with N, O or S, and wherein the optional substituents are selected from carbonyl, carboxy, hydroxy, amino and other groups, and Y is the covalent linkage between A and Z where Y is selected from -NHCO-, -CONH-, -CONHNHCO-, -N=N-, -N=CH-, -NHCH₂-, -NHN=CH-, -NHNHCH₂-, -SS-, -SCH₂-, -CH₂S-, -NHCRNH-, -COO-, -OCO-, and R is O, S or NH₂, and Z is GCSF or EPO. The invention also describes reagents that can be used as probes for the detection of buried thiol groups of a protein or peptide, said reagent comprising a complex of either vitamin B₁₂ (or an analogue thereof) or more generally of any instrumentally or visually detectable label, covalently linked to a diradical spacer, said spacer having a terminal reactive group capable of forming a disulphide bond with a free thiol in said protein or peptides.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

VITAMIN B₁₂ MEDIATED ORAL DELIVERY SYSTEMS FOR GCSF AND EPO

5 BACKGROUND OF THE INVENTION

10 The present invention relates to the oral delivery of the therapeutic substances granulocyte-colony stimulating factor (GCSF) and erythropoietin (EPO) by administration of a complex comprising these substances linked to vitamin B₁₂ (VB₁₂) or an analogue thereof. More particularly, the invention relates to methods for the synthesis of these complexes and to methods for the amplification of the amount of GCSF or EPO delivered per VB₁₂ carrier molecule.

15 An oral delivery system is known, because of recent work undertaken by one of the current inventors, which is described in PCT Application WO87/02251 (PCT/AU86/0299), whereby an active substance linked to at least one carrier molecule, which is VB₁₂ or an analogue thereof, can use the natural VB₁₂ uptake system mediated by the binding of VB₁₂ to intrinsic factor (IF) to transport the resultant complex from the intestinal lumen into the circulation. Once delivered into serum or the lymphatic drainage system the complex substantially
20 retains the bioactivity of the native active substance.

In common with virtually all proteins, peptides and other large bioactive molecules there is currently no method for the oral delivery of either GCSF or EPO. The oral route of administration is the most preferable means of delivering a pharmaceutically active agent,
25 and as such there is a large and valuable market for any process which permits the oral delivery of either of these proteins to humans. Such a process would be available by the formation of a complex between VB₁₂ and GCSF or EPO.

SUMMARY OF THE INVENTION

30 The present invention relates in one aspect to complexes between VB₁₂ analogues and either GCSF or EPO that retain both significant affinity for intrinsic factor (IF) in the VB₁₂

portion of the complex and significant bioactivity of the GCSF or EPO portion of the complex. The invention also concerns a process for the synthesis of these complexes. This may be achieved at least in part, by using a spacer compound, which is linked covalantly
5 between the VB₁₂ portion and the GCSF or EPO.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

One aspect of the invention provides a complex which comprises at least one active substance linked to at least one carrier molecule, which is VB₁₂ or an analogue of VB₁₂
10 wherein the ability of the carrier to undergo the binding reactions necessary for uptake and transport of VB₁₂ in a vertebrate host and the activity of the active substance are substantially maintained. This occurs by providing a complex between GCSF and a carrier selected from vitamin B₁₂ or a vitamin B₁₂ analogue wherein the GCSF and the carrier are covalently linked through a diradical spacer, the complex being capable of binding to
15 intrinsic factor with high affinity with maintenance of GCSF bioactivity.

The complex preferably has the general form V-X-A-Y-Z, where V is the carrier molecule VB₁₂ or an analogue (including derivatives) thereof that retains IF affinity, Z is the active substance selected from EPO or GCSF, A is a spacer arm of variable composition and length, X is the functional group through which V is attached to A, and Y is the
20 functional group through which Z is attached to A. The nature of functional group X, its site of attachment to V and the nature of spacer arm A are chosen to maximise the IF affinity of the complex. The nature of functional group Y, its site of attachment to Z and the nature of spacer arm A are chosen to maintain substantially the bioactivity of Z.

25 Preferably X is selected from: -NHNH-, -NH-, -O-, S-, -SS- or -CH₂-. A is preferably an optionally substituted, saturated or unsaturated, branched or linear, C₁₋₅₀ alkylene, cycloalkylene or aromatic group, optionally with one or more carbons within the linear chain being replaced with N, O or S, and wherein the optional substituents are selected
30 from carbonyl, carboxy, hydroxy, amino and other groups. Y is preferably a covalent linkage between spacer chain A and protein Z, where Y is selected from -NHCO-, -

CONH-, -CONHNHCO-, -N=N-, -N=CH-, NHCH₂-, -NHN=CH-, -NHNHCH₂-, -SS-, -SCH₂-, -CH₂S-, -NHCRNH- [where R = O, S or NH₂], -COO-, -OCO-, and Z is GCSF or EPO.

5

In the context of the present invention, the term "active substance" (ie Z) includes all, part or an analogue, homologue, derivative or combination thereof of either granulocyte colony stimulating factor (GCSF) or erythropoietin (EPO).

10

The carrier is VB₁₂ or VB₁₂ analogue. The VB₁₂ analogues include any variant or derivative of VB₁₂ (cyanocobalamin) which possesses binding activity to intrinsic factor. Preferred analogues of VB₁₂ also include aquocobalamin, adenosylcobalamin, methylcobalamin, hydroxycobalamin, cyanocobalamin, carbanalide, and 5-methoxybenzylcyanocobalamin [(5-MeO)CN-Cbl] as well as the desdimethyl, monoethylamide and the methylamide analogues of all of the above. Other analogues include all alkyl cobalamins in which the alkyl chain is linked to the corrin nucleus by a direct CoC covalent bond. Other analogues include chlorocobalamin, sulfitocobalamin, nitrocobalamin, thiocyanatocobalamin, benzimidazolecyanocobalamin derivatives such as the: 5,6-dichlorobenzimidazole, 5-hydroxybenzimidazole, trimethylbenzimidazole, as well as adenosylcyanocobalamin [(Ade)CN-Cbl], cobalamin lactone, cobalamin lactam and the anilide, ethylamide, monocarboxylic and dicarboxylic acid derivatives of VB₁₂ or its analogues.

15

20

25

Preferred derivatives of VB₁₂ also include the mono-, di- and tricarboxylic acid derivatives or the propionamide derivatives of VB₁₂. Carriers may also include analogues of VB₁₂ in which the cobalt is replaced by zinc or nickel. The corrin ring of VB₁₂ or its analogues may also be substituted with any substituent which does not effect its binding to IF, and such derivatives of VB₁₂ or its analogues are part of this invention. Other derivatives of VB₁₂ or its analogues which have a functional group which is able to react with the spacer compound are also part of the invention.

30

It is preferred that the complex may comprise GCSF linked through a disulphide bond to

a (dithiopyridyl propionamido) dodecylamino [DTP-dodecylamino] derivative of VB₁₂.

5 Another preferred embodiment of the invention provides a process for the production of a complex comprising GCSF linked through a disulphide bond to a (dithiopyridyl propionamido) dodecylsuberylhexylamino derivative of VB₁₂, (a long-chain analogue of the DTP-dodecylamino VB₁₂) which displays higher affinity for intrinsic factor.

10 Another preferred embodiment of the invention provides a process for the production of a complex comprising GCSF linked through a disulphide bond to an (dithiopyridyl propionamido) dodecylcarboxamidomethyl derivative of VB₁₂, in which the spacer is linked to the VB₁₂ through an axial CoC bond.

15 Another embodiment of the invention provides a process for the production of a complex comprising at least one active substance linked to at least one carrier molecule through a spacer, the carrier molecule being VB₁₂ or an analogue thereof, wherein the ability of the carrier to undergo the binding reactions necessary for uptake and transport of VB₁₂ in a vertebrate host and the activity of the active substance are substantially maintained, the process comprising one or more of the following steps:

- 20 a) reacting together the active substance and the carrier and the spacer compound to form the complex;
- b) reacting the active substance with the spacer, and then reacting the product with the carrier to form the complex;
- 25 c) reacting the carrier with the spacer, and then reacting the product with the active substance to form the complex;
- d) following method of (a), (b) or (c) but with the additional step of having chemically modified the carrier and/or the active substance in a previous step to provide a functional group on the carrier and/or active substance which will react with the spacer compound; or
- 30 e) following the method of (a), (b), (c) or (d) but with the additional step of reacting the active substance or the carrier with a polymeric support, before carrying out the

further reactions. Preferably the polymeric support is bonded both to the carrier (or spacer-carrier), and also to the active substance or (spacer-active substance).

5 For example, it is possible to form these complexes using the "e" mono -acid of "e"VB₁₂ by: (i) preparing the mono-acid derivative of VB₁₂ by mild acid hydrolysis of cyanocobalamin and purifying the initial hydrolysate; (ii) modifying the e-mono-acid to give a terminal functional group attached to the eVB₁₂ nucleus through a spacer arm; (iii) coupling the functionalised eVB₁₂ derivative to carboxylate, amine, thiol, hydroxyl, phenol,
10 aldehyde or ketone groups or other suitable functional groups present initially or introduced chemically on the active substance. The spacer compound can be selected to have suitable functional groups at either end of its backbone, or else these functional groups can be introduced, if necessary by normal chemical synthetic reactions.

15 The invention also involves the modification of a polymeric support to introduce functional groups capable of reacting either directly with the spacer compound or with the spacer linked with the active substance. The resulting polymer-active substance intermediate ideally contains many molecules of the active substance, and this intermediate is suitable for coupling to the carrier to give a complex capable of amplified delivery of the active
20 substance.

The invention also concerns a general method for the modification of unreactive thiols in peptides and proteins, particularly those not normally exposed to reagents dissolved in aqueous solvents because they are buried in hydrophobic regions of the protein. These
25 modified peptides and proteins can then be labelled, if desired.

The invention also involves a reagent , of the general formula:



wherein, V' is vitamin B₁₂ or a vitamin B₁₂ analogue, bonded to X either through a
30 carboxylate group pendant to the corrin nucleus of VB₁₂ or through the central cobalt atom or to a functional group introduced onto the VB₁₂ molecule, or V' is a label, where "label"

refers to any substance that is detectable by visual or instrumental methods. Labels can include chromogens, catalysts, fluorescent compounds, chemiluminescent compounds, radioactive isotopes, colloidal metal and non-metallic particles, dye particles, enzymes or substrates, antibodies or antigens, biotin, avidin or streptavidin, latex particles, liposomes or other vesicles containing signal producing substances, and the like.

X is selected from -NHCO-, -CONH-, -CONHNHCO-, -N=N-, -N=CH-, -NHCH₂-, -NHN=CH-, -NHNHCH₂-, -SS-, -SCH₂-, CH₂S-, -NHCRNH-, [R is O, S or NH₂], -COO-, -OCO-, and A is an optionally substituted, saturated or unsaturated, branched or linear, C₁₋₅₀ alkylene, cycloalkylene or aromatic group, optionally with one or more carbons within the chain being replaced with N, O or S, and wherein the optional substituents are selected from carbonyl, carboxy, hydroxy, amino and other groups, and Y' is a functional group capable of reacting with thiols to give a stable covalent linkage, including iodoacetyl, bromoacetyl, chloroacetyl, maleimido, 3-carboxy-4-nitrophenyldithio or 2-pyridyldithio, and similar groups.

The complexes described herein may be formulated into pharmaceutical or veterinarily acceptable compositions utilising carriers and/or excipients as are well known in the art. Examples of suitable dosage forms which may be used in this invention are described for example in *Remington's Pharmaceutical Sciences (Mack Publishing Company, 10th Edition, which is incorporated herein by reference)*. Compositions may be in the form of a capsule, tablet, slow-release dosage form, elixir, gel, paste, enterically coated dosage form, or any other suitable dosage form as is well known in the art.

Complexes and compositions according to this invention may be administered to a human or animal subject in need of treatment with GCSF or EPO. Modes of administration are not critical to this invention and include parenteral (intravenous, intramuscular, or intraorgan injection), oral, transdermal, vaginal, anal, or other administration routes as are well known in the art. A therapeutically effective amount of a complex or compound according to this invention is that which provides treatment of a particular disease state.

What constitutes an effective amount will depend upon the nature of the disease being treated, the consulting physician or veterinary surgeon judgement, and other factors such as the age, weight and or sex of the subject. By way of example only, an effective amount of a complex composition according to the invention may comprise from one nanogram to 10 grams of a complex in accordance with the invention.

This invention also relates to the use of complexes described herein for the manufacture of medicament, and for the treatment of conditions responsive to GCSF or EPO.

BEST METHOD OF CARRYING OUT THE INVENTION

EXAMPLES

Materials: VB₁₂ was obtained from Rousell-Uclaf. GCSF and EPO were obtained from Amgen. 1-Ethyl-3-(dimethylaminopropyl) carbodiimide.HCl (EDAC.HCl) was obtained from Biorad. N-Succinimidyl 3-(2-pyridyldithio) propionate (SPDP) and succinimidyl, 6-[3-(2-pyridyldithio) propionamido] hexanoate (LC-SPDP) were obtained from Pierce Chemical Co. All other reagents were obtained from Fluka.

Example 1: VB₁₂ -GCSF complexes

Three classes of VB₁₂-GCSF complexes were prepared:

- a) conjugated via an amide linkage, formed by carbodiimide (EDAC) mediated coupling of an amino terminal eVB₁₂ derivative to the C-terminus of GCSF or the carboxylate side chains of GCSF.
- b) conjugated via a disulphide linkage formed by a thiol insertion reaction of the free thiol at Cys-17 of GCSF into the disulphide bond present in the (dithiopyridylpropionamido) terminal derivatives of eVB₁₂.
- c) conjugated via an acyl hydrazide linkage, formed by EDAC-mediated coupling of a hydrazido-terminal eVB₁₂ derivative to the C-terminus of GCSF or the carboxylate side chains of GCSF.

1.1 Production and purification of the "e" isomer of monocarboxy-VB₁₂.

The "e" isomer of monocarboxy vitamin B₁₂, formerly named the d isomer, but reassigned by Anton and co-workers (1980: J. Am. Chem. Soc. 102:2215) as the e isomer, was separated from the b and d isomers formed during acid hydrolysis of cyanocobalamin by a combination of Dowex 1X2 chromatography and semi-preparative C-18 RP-HPLC developed with a gradient of acetonitrile in 0.1% TFA.

1.2 Production of amino-derivatives of eVB₁₂

A number of amino-derivatives of eVB₁₂ were prepared by reacting the e isomer with:

- i) 1,2-diaminoethane
- ii) 1,6-diaminohexane
- iii) 1,12-diaminododecane
- iv) 1,3-diamino-2-hydroxypropane
- v) 1,6-diamino-3,4-dithiahexane (a.k.a. cystamine)

All reactions were performed at pH 6.5 using a twenty fold molar excess of the diamine over e isomer and a twenty fold molar excess of EDAC. In a typical reaction 135 mg of eVB₁₂ was dissolved in distilled water (6 ml) to which was added 1.2 ml of 1.0 M diamine, pH 6.5. Dry EDAC (270 mg) was then added and the reaction mixture was left overnight at room temperature.

All amino derivatives were purified by reverse phase chromatography on a semi-preparative C-4 column using a 5-100% acetonitrile gradient in 0.1% TFA. Eluted material was further purified by S-Sepharose chromatography. The amino-derivative was eluted with 0.1 M HCl, followed by extraction into phenol, and back-extraction into water after the addition of dichloromethane to the phenol phase. The amino-eVB₁₂ derivatives were then recovered from the water phase by lyophilization.

1.3 Conjugation of 2-aminoethyl-eVB₁₂ to GCSF

A solution of 2-aminoethyl-eVB₁₂ (26.5 mg, 18 μ mol) in 2 ml of GCSF (6 mg/ml, 0.63 μ mol) was cooled to 4 C. An aliquot of freshly prepared EDAC solution (100 mg/ml, 120

μl , 63 μmol) was added. After 24 h at 4 C a second aliquot of freshly prepared EDAC solution was added. The reaction was allowed to proceed for a total of 48 h at 4 C, after which the unreacted 2-aminoethyl-eVB₁₂ was separated from the conjugate and aggregate by chromatography on Sephadex G-50 in 2.5% acetic acid.

1.4 Conjugation of cystaminyI-eVB₁₂ to GCSF

A solution of cystaminyI-eVB₁₂ hydrochloride (30 mg, 20 μmol) in 2.5 ml of GCSF (6 mg/ml, 0.80 μmol) at 4 C was treated with an aliquot of an aqueous EDAC solution (20 mg/ml, 75 μl , 8 μmol). Further aliquots of freshly prepared EDAC solution (20 mg/ml, 75 μl , 8 μmol) were added after 4.5 h, 7 h and 24 h. The reaction was allowed to proceed for a total of 48 h at 48 C, after which time the unreacted amino-eVB₁₂ was separated from the conjugate by chromatography on Sephadex G-50 in 2.5% acetic acid.

1.5 Preparation of 3-(2-pyridyldithio) propionamido derivatives of aminoethyl-eVB₁₂

The dithiopyridyl derivatives of amino-eVB₁₂ were prepared by reacting SPDP with:

- i) 2-aminoethyl-eVB₁₂
- ii) 6-aminoethyl-eVB₁₂
- iii) 12-aminododecyl-eVB₁₂

In a typical reaction the terminal amino-eVB₁₂ was dissolved at 50 mg/ml in 0.1 M PO₄ buffer, pH 7.5, containing 0.1 M NaCl. SPDP was dissolved at 50 mg/ml in acetone and 800 μl of the solution was added to the amino-eVB₁₂. After reaction overnight at room temperature the DTP-amino-eVB₁₂ was purified by RP-HPLC on a semi-prep C4 column, and then lyophilized.

1.6 Conjugation of DTP-amino-VB₁₂ to GCSF

In initial conversations with AMGEN it was revealed that they had found that it was impossible to modify the free cysteine in undenatured GCSF with standard thiol modifying agents. Initial experiments with DTP-aminoethyl-eVB₁₂ showed that it was possible to achieve some 20% substitution of GCSF with the VB₁₂ in the absence of guanidine; this

level rose to >80% in the presence of 4 M guanidine. It was therefore decided that it might be possible to access the free thiol with DTP-amino-eVB₁₂ in the absence of guanidine if a longer spacer was used for the conjugation.

5

In a second series of experiments GCSF was reacted with DTP-aminoethyl-, DTP-aminohexyl- and DTP aminododecyl-eVB₁₂ in the presence or absence of 4 M guanidine in 0.1 M sodium acetate buffer, pH 4.0.

10 The degree of substitution of GCSF by various DTP-amino-eVB₁₂ -spacer complexes is shown in the following table:

TABLE 1

	<u>Spacer</u>	<u>Guanidine</u>	<u>+ Guanidine</u>
	DTP-aminoethyl-	37.5%	89.3%
15	DTP-aminohexyl-	45.5%	95.2%
	DTP-aminododecyl-	100.0%	100.0%

From Table 1 it can be seen that by switching to the longer dodecyl-spacer it was possible to conjugate to the buried thiol in GCSF without the use of guanidine.

20

The initial attempts to conjugate to the free thiol group in GCSF using the pyridyldithiopropionamido-aminoethyl derivative of eVB₁₂ resulted in a small degree of conjugation, of around 20-40 percent in the absence of guanidine. The addition of 4 M guanidine (final concentration) raised the conjugation efficiency to over 80%. Preparation of a longer, more hydrophobic derivative of eVB₁₂, pyridyldithiopropionamido-dodecyl eVB₁₂ resulted in 100% substitution of GCSF after 24 h at 4 C, without the need for the addition of guanidine. The use of the thiol interchange chemistry in this reaction proved advantageous as the eVB₁₂ conjugation was surprisingly successful at pH's which minimised the extent to which GCSF undergoes spontaneous aggregation. Chromatography of the conjugated material resulted in base-line separation of conjugate from free eVB₁₂.

30

This type of approach can be more generally applied for the development of reagents capable of detecting, quantifying and/or modifying thiol groups in proteins and peptides which were

hitherto regarded as chemically unreactive.

1.7 Scale up conjugation of DTP-aminododecyl-eVB₁₂ to GCSF

Following the initial success with the conjugation of the DTP-dodecyl-spacer, the reaction was scaled up as follows:

To 2.5 ml of GCSF (6 mg/ml; 15 mg) was added 1.6 ml of DTP-aminododecyl-eVB₁₂ (10 mg/ml in 2.5% acetic acid). The reaction was allowed to proceed for 48 h at 4 C, after which the unreacted eVB₁₂ was separated from the conjugate by chromatography on Sephadex G-25 in 2.5% acetic acid. Fractions containing GCSF were pooled, concentrated in an AMICON positive pressure cell using a YM 10 membrane and dialysed for 72 h against sterile distilled water.


1.8 Preparation of DTP-dodecylsuberlhexyl-eVB₁₂ reagent.

Although the DTP-dodecyl-eVB₁₂ reagent reacts efficiently with GCSF to give a stable, well characterised complex, the material had a low IF affinity (~2-3% of native eVB₁₂). A complex with increased IF affinity was prepared by synthesising an extended spacer analogue of the DTP-dodecylamino-eVB₁₂. The synthesis of this analogue uses the same SPDP chemistry to conjugate through the cysteine of GCSF, however a longer spacer arm is attached to the eVB₁₂. As anticipated, this resulted in the formation of a conjugate whose IF affinity is significantly greater than that of the conjugate prepared by the reaction of GCSF with DTP-dodecyl-eVB₁₂ (see Table 2).

This spacer was prepared by sequential reaction of the e-carboxylate of eVB₁₂ with:

- i) 1,6-diaminohexane and EDAC (to give 6-aminohexyl eVB₁₂).
- ii) disuccinimidylsuberate (DSS) (to give monosuccinimidylsuberylhexyl-eVB₁₂).
- iii) 1,12-diaminododecane (to give 12-aminododecylsuberylhexyl-eVB₁₂).

That is:



The resultant spacer, which is more than twice the length of aminododecyl-eVB₁₂, was derivatized at the terminal amino group with SPDP and coupled to GCSF by means of the protocol described in the following section.

1.9 Conjugation of DTP-dodecylsuberylhexyl-eVB₁₂ to GCSF

A solution of the DTP-dodecylsuberylhexyl-eVB₁₂ (9 mg, 7 μmol) was taken up in acetic acid (100 μl) and diluted to 1 ml with water. This solution was added to 5 ml of GCSF solution (4 mg/ml, 0.5 μmol) cooled to 4 °C. The reaction mixture was left for 144 h at 4 °C, then worked up using the standard protocol.

1.10 Production of hydrazide derivatives of eVB₁₂ carboxylate


Two hydrazide derivatives of eVB₁₂ carboxylate were prepared for conjugation to carboxyl groups of GCSF by reaction with EDAC. The two hydrazide derivatives used, and their (shorthand) chemical structure, are:

- a) hydrazido-eVB₁₂ (= eVB₁₂-CONHNH₂)
- (b) adipyl-hydrazido-eVB₁₂ (= eVB₁₂-CONHNHCO(CH₂)₄CONHNH₂)

1.10i Hydrazido-eVB₁₂


This reagent was prepared by a two step synthesis involving the coupling of tert butyl carbazate to eVB₁₂ carboxylate and subsequent removal of the tBoc group to generate the

free hydrazide.




1.10ii Adipyl-hydrazido-eVB₁₂

This reagent was readily prepared in one step from eVB₁₂ carboxylate by the addition of EDAC to a mixture of the acid and a 20-fold excess of adipylhydrazide.



EDAC-mediated coupling of hydrazido-eVB₁₂ analogues to the carboxylate side-chains of GCSF proceeded more readily, and required significantly lower amounts of eVB₁₂ derivative and EDAC, than the corresponding conjugations of amino eVB₁₂ derivatives to GCSF. This is readily explainable in terms of the relative basicity of hydrazides (pK_a 2.6) in comparison with amines (pK_a 8-9). Thus at the pH at which the GCSF coupling takes place (4-5) a hydrazido eVB₁₂ derivative will be primarily in the reactive, non-protonated form while an amino-eVB₁₂ derivative will be primarily in the non-reactive, protonated form.

The coupling reaction used is:



1.11 Conjugation of hydrazido-eVB₁₂ to GCSF

A solution of hydrazido eVB₁₂ (8.9 mg, 6.5 μ mol) in 5 ml of GCSF solution (4 mg/ml, 1.05 μ mol) was cooled to 4 C and an aliquot of EDAC solution (50 mg/ml, 40 μ l, 10 μ mol) was added. After 5 h an identical aliquot of fresh EDAC solution was added and the reaction mixture was left overnight at 4 C. Conjugate was removed from unreacted eVB₁₂ and other reagents by chromatography on Sephadex G-50 in 2.5% acetic acid.

1.12 Conjugation of adipylhydrazido-eVB₁₂ to GCSF

A solution of adipylhydrazido eVB₁₂ (10 mg, 6.6 μ mol) in 3 ml of GCSF solution (4 mg/ml, 0.64 μ mol) was cooled to 4 C and an aliquot of EDAC solution (50 mg/ml, 40 μ l, 10 μ mol) was added. After 5 h a 20 μ l aliquot of fresh EDAC solution (50 mg/ml, 20 μ ml, 5 μ mol) was added and the reaction mixture was left overnight at 4 C. Conjugate was removed from unreacted eVB₁₂ and other reagents by chromatography on Sephadex G-50 in 2.5% acetic acid.

1.13 Preparation of poly-GCSF-HPMA-eVB₁₂ complex.

Two N-(2-hydroxypropyl)methacrylamide (HPMA) copolymers can be synthesized as polymer backbones for the incorporation and derivatization with GCSF and eVB₁₂. A non-biodegradable polymer backbone (HPMA-GG) can be synthesized by the free radical copolymerization of HPMA with N-methacryloylglycylglycine p-nitrophenyl ester. A biodegradable polymer (HPMA-FALG) can be synthesized by the free radical copolymerization of HPMA with N-methacryloylglycylphenylalanylleucylglycine p-nitrophenol ester by the method of Rejmanova and coworkers [Rejmanova, P, Obereigner, B and Kopecek, J, 1981 Makromol Chem 182:1899-1915]. In order to incorporate GCSF and eVB₁₂ onto the polymers, the polymers are initially reacted with a ten molar excess of a mixture of aminododecyl-eVB₁₂ and dithiopyridyldodecylsuberylhexylamine (1:10 mole:mole) overnight. Unreacted nitrophenyl esters are subjected to aminolysis by the addition of 1-amino-2-propanol. The modified polymers are purified by chromatography on Sepharose 6B. A solution of the dithiopyridyldodecylsuberylhexyl modified eVB₁₂-substituted polymers is dissolved in 2.5% acetic acid and reacted with GCSF. The reaction mixture is left for 144 h at 4 C, after which the GCSFeVB₁₂-substituted polymers can be

purified by chromatography on Sepharose 6B.

Part 2: eVB₁₂-EPO complexes

5 Three classes of eVB₁₂-EPO complexes were prepared:

- (a) Conjugated via an amide linkage, formed by EDAC-mediated coupling of an amino eVB₁₂ derivative to the C-terminus of EPO, the carboxylate side chains of the Asp/Glu residues of EPO or the carboxylate groups of the sialic acid residues of the carbohydrate portion of EPO.
- 10 (b) Conjugated via an acyl hydrazide linkage, formed by EDAC-mediated coupling of a hydrazido-eVB₁₂ derivative to the carboxylate side chains of the Asp/Glu residues of EPO or the carboxylate groups of the sialic acid residues of the carbohydrate portion of EPO.
- (c) Conjugated via a hydrazone linkage between a hydrazido-eVB₁₂ derivative and an
15 aldehyde group generated by periodate oxidation of the carbohydrate residues of EPO.

2.1 Conjugation of 2-aminoethyl-eVB₁₂ to EPO

20 A mixture of 2-aminoethyl-eVB₁₂ (8 mg, 5.7 μ mol) and EPO (27 mg/ml, 200 μ l, 0.18 μ mol) was cooled to 4 C and an aliquot of EDAC solution (10 mg/ml, 100 μ l, 5 μ mol) was added. The reaction mixture was left for 64 h at 4 C and finally purified by size-exclusion chromatography on a Superdex-75 column. Elution with a buffer consisting of Tris (pH 7.5, 10 mM; NaCl, 100 mM) afforded the purified EPO-eVB₁₂ complex.

25 2.2 Conjugation of cystaminyl-eVB₁₂ to EPO

A mixture of 6-amino-3,4-dithiahexylamide-eVB₁₂ (12 mg, 8.1 μ mol) and EPO (13.5 mg/ml, 500 μ l, 0.23 μ mol) was cooled to 4 C and an aliquot of EDAC solution (10 mg/ml, 250 μ l, 13 μ mol) was added. The reaction mixture was left for 48 h at 4 C and was finally purified by size-exclusion chromatography on a Sephadex G-75 column. Elution with a
30 buffer consisting of Tris (pH 7.5, 10 mM)/NaCl (100 mM) afforded the purified EPO-eVB₁₂ complex.

2.3 EDAC-mediated conjugation of hydrazido-eVB₁₂ to EPO

A solution of hydrazido VB₁₂ (10 mg, 7.3 μ mol) in 7 ml of EPO solution (2.6 mg/ml, 0.6 μ mol) was cooled to 4 C and an aliquot of EDAC solution (20 mg/ml, 50 μ l, 5 μ mol) was added. After 5 h a second aliquot of fresh EDAC solution (10 mg/ml, 25 μ l, 1.3 μ mol) was added and the reaction mixture was left overnight at 4 C. The conjugate was purified by size-exclusion chromatography on a G-50 column. Elution with a buffer consisting of Tris (pH 7.5, 10 mM; NaCl, 100 mM) afforded the purified EPO-eVB₁₂ complex .

2.4 EDAC-mediated conjugation of adipyl-hydrazido-eVB₁₂ to EPO

A solution of adipyl-hydrazido-eVB₁₂ (11 mg, 7.3 μ mol) in 4 ml of EPO solution (2.6 mg/ml, 0.35 μ mol) was cooled to 4 C and an aliquot of EDAC solution (10 mg/ml, 100 μ l, 5 μ mol) was added. The reaction mixture was left overnight at 4 C. The conjugate was purified by size-exclusion chromatography on a G-50 column. Elution with a buffer consisting of Tris (pH 7.5, 10 mM)/NaCl (100 mM) afforded the purified EPO-eVB₁₂ complex .

2.5 Periodate mediated conjugation of hydrazido-eVB₁₂ to EPO

An aqueous solution of EPO (6.4 mg/ml, 1.56 ml, 0.33 μ mol) was cooled to 4 C and freshly prepared sodium periodate solution (25 mM, 360 μ l) was added. The solution was left to stir gently for fifteen minutes at 4 C and the excess periodate was quenched by the addition of ethylene glycol (5 μ l). The solution was dialyzed overnight against 3 l of pH 5.6, 10 mM NaOAc buffer. A solution of hydrazido-eVB₁₂ (7.5 mg, 5 μ mol) in distilled water (200 μ l) was added to the oxidised EPO solution and the reaction mixture was left for 7 h at 4 C. The conjugate was separated from unreacted hydrazide by SEC on G-75 Sephadex. Elution with a buffer consisting of Tris (pH 7.5/10 mM)/NaCl (100 mM) afforded the purified EPO-eVB₁₂ complex .

2.6 Preparation of poly-EPO-HPMA-eVB₁₂

Two N-(2-hydroxypropyl)methacrylamide (HPMA) copolymers can be synthesized as polymer backbones for the incorporation and derivatization with EPO and VB₁₂. A

nonbiodegradable polymer backbone (HPMA-GG) is synthesized by the free radical copolymerization of HPMA with N-methacryloylglycylglycine p-nitropheny ester. A biodegradable polymer (HPMA-GFLAG) is synthesized by the free radical copolymerization of HPMA with N-methacryloylglycyl-phenylalanylleucylglycine p-nitrophenol ester by the method of Rejmanova and coworkers [Rejmanova, P, Obereigner, B and Kopecek J, 1981 Makromol Chem 182:1899-1915]. In order to incorporate EPO and eVB₁₂ onto the polymers, the polymers are reacted with a ten molar excess of a mixture of aminododecyl-eVB₁₂ and 6-(3-acylhydrazidylpropionamido)-1-amino-3,4-dithiahexane (1:10 mole:mole) overnight. Unreacted nitrophenyl esters are subjected to aminolysis by the addition of 1-amino-2-propanol. The modified polymers are purified by chromatography on Sepharose 6B. EPO is covalently linked to the 6-(3-acylhydrazidylpropionamido)-1-amino-3,4-dithiahexane modified eVB₁₂-substituted polymers by the addition of EDAC. The reaction mixture is left for 18 h at 4 C, after which the EPO-eVB₁₂-substituted polymers are purified by chromatography on Sepharose 6B.

Example 3:

The diradical spacer may be selected with a suitable length and functionality to optimize the intrinsic factor (IF) affinity of the resulting GCSF or EPO eVB₁₂ complex. As examples of this, the following table shows the result of altering the length and functionality of the spacer for various complexes and VB₁₂ derivatives.

The IF affinity of the various eVB₁₂ analogues and complexes referred to in the Examples are shown in Table 2. In particular, there is significant increase in IF affinity between the GCSF-dithiopropionamido[12-aminododecylamido]-eVB₁₂ (prepared in section 1.7) and the GCSF- dithiopropionamido[12-aminododecyl-subeyl-hexylamido]-eVB₁₂ (prepared in section 1.9). This increase in IF affinity presumably results from the increase in length of the spacer arm in the latter complex.

Table 2

Relative Intrinsic Factor affinity of VB₁₂ analogues and complexes

	<u>eVB₁₂ Analogue or complex</u>	<u>IF Affinity</u>
	eVB ₁₂ carboxylate	35%
5	2-aminoethylamido-eVB ₁₂	48%
	6-aminohexylamido-eVB ₁₂	91%
	12-aminododecylamido-eVB ₁₂	74%
	12-aminododecyl-suberyl-hexylamido-eVB ₁₂	82%
10	dithiopropionamido[2-aminoethylamido]-eVB ₁₂	6%
	dithiopropionamido[6-aminohexylamido]-eVB ₁₂	7%
	dithiopropionamido[6-aminohexylamido]-eVB ₁₂	11%
	GCSF-dithiopropionamido[12-aminododecylamido]-eVB ₁₂	3%
15	GCSF-dithiopropionamido12-aminododecyl-suberyl-hexylamido-eVB ₁₂	28%
	hydrazidyl-eVB ₁₂	100%
	adipyl dihydrazidyl-eVB ₁₂	44%
	GCSF-acylhydrazidyl-eVB ₁₂	10%
20	GCSF-acyl[adipyl dihydrazidyl]-eVB ₁₂	7%
	EPO-acylhydrazidyl-eVB ₁₂	6%
	EPO-acyl[adipyl dihydrazidyl]-eVB ₁₂	11%
25	IF Affinity is expressed as a percentage of the affinity of the eVB ₁₂ derivative relative to native, unmodified VB ₁₂ .	

THE CLAIMS:

1. A complex between GCSF and a carrier comprising vitamin B₁₂ or a vitamin B₁₂ analogue wherein said GCSF and said carrier are covalently linked through a diradical spacer, said complex being capable of binding to intrinsic factor with high affinity with maintenance of GCSF bioactivity.

2. A complex according to claim 1 wherein said carrier is covalently linked to GCSF through: a disulphide bond with Cys-17 of GCSF; an amide linkage; an acyl hydrazide linkage; an imine linkage; or a hydrazone linkage.

3. A complex according to claim 1, of the general formula:



wherein,

V is vitamin B₁₂ or a vitamin B₁₂ analogue, or derivative, bonded to X either through a carboxylate group pendant to the corrin nucleus of VB₁₂ or through the central cobalt atom or to a functional group introduced onto the VB₁₂ molecule,

X is selected from: -NHNH-, -NH-, -O-, S-, -SS- or -CH₂-, and

A is an optionally substituted, saturated or unsaturated, branched or linear, C₁₋₅₀ alkylene, cycloalkylene or aromatic group, optionally with one or more carbons within the linear chain being replaced with N, O or S, and wherein the optional substituents are selected from carbonyl, carboxy, hydroxy, amino and other groups, and

Y is the covalent linkage between A and Z where Y is selected from -NHCO-,

-CONH-, -CONHNHCO-, -N=N-, -N=CH-, -NHCH₂-, -NHN=CH-, -NHNHCH₂-, -SS-, -SCH₂-, -CH₂S-, -NHCRNH-, [R is O, S or NH₂], -COO-, -OCO-, and , Z is GCSF.

4. A complex according to claim 1 wherein said vitamin B₁₂ analogue is selected from cyano-cobalamin (CN-Cbl), aquocobalamin, adenosylcobalamin, methylcobalamin,

hydroxy-cobalamin, (5-methoxybenzyl)cyanocobalamin [(5-MeOBz)CN-Cbl]; the desdimethyl, monoethylamide and the methylamide analogues of these; alkyl cobalamins in which the alkyl chain is linked to the corrin nucleus by a direct Co C covalent bond; chlorocobalamin, sulfitocobalamin, nitrocobalamin, thiocyanatocobalamin, benzimidazole derivatives including 5,6-dichlorobenzimidazole, 5-hydroxybenzimidazole, trimethylbenzimidazole; adenosylcyanocobalamin [(Ade)CN-Cbl], cobalamin lactone, cobalamin lactam; the anilide, ethylamide, monocarboxylic and dicarboxylic acid derivatives of VB₁₂ or its analogues; the mono-, di- and tricarboxylic acid derivatives of VB₁₂; and analogues of VB₁₂ in which the cobalt is replaced by zinc or nickel.

5. A complex according to claim 1 wherein said vitamin B₁₂ analogue is an analogue where the corrin ring is substituted with a substituent which does not affect binding to intrinsic factor.

6. A complex according to claim 1 wherein said GCSF is covalently bound to a pharmaceutically acceptable polymer.

7. The complex according to claim 6 wherein the polymer is selected from dextran, inulin, cellulose, starch and derivatives thereof, chondroitin sulfate, poly[N- α -(2-hydroxypropyl)-methacrylamide] and derivatives thereof, styrene-maleic anhydride copolymer, divinylether-maleic anhydride copolymer, polylysine, poly(glutamic acid), poly(hydroxypropyl glutamine), poly(lactic acid), water soluble polyurethanes formed by covalent linkage of PEG with lysine or other amino acids and branched chain polypeptides.

8. A complex according to claim 6 wherein said pharmaceutically acceptable polymer is biodegradable within the human or animal body.

9. A process for the production of a complex between GCSF and a carrier selected from

vitamin B₁₂ or a vitamin B₁₂ analogue which comprises: reacting one or both of said carrier and GCSF with a diradical spacer having terminal reactive groups to form a carrier/linker and/or GCSF linker intermediate, and thereafter reacting together the
5 respective components to give a complex between GCSF and said carrier wherein the components are covalently linked through said spacer.

10 10. A process according to claim 9 wherein GCSF and/or said carrier are modified prior to complex formation to provide at least one functional group capable of forming a chemical linkage.

15 11. A process according to claim 9 wherein said carrier is covalently linked to GCSF through a disulphide linkage with Cys-17 of GCSF, an amide linkage, an acyl hydrazide, an imine linkage, or a hydrazone linkage.

20 12. A complex between EPO and a carrier comprising vitamin B₁₂ or a vitamin B₁₂ analogue wherein said EPO and said carrier are covalently linked through a diradical spacer, said complex being capable of binding to intrinsic factor with high affinity with maintenance of EPO bioactivity.

25 13. A complex according to claim 12 wherein said carrier is covalently linked to EPO through: an amide linkage; an acyl hydrazide linkage; an imine linkage; or a hydrazone linkage.

30 14. A complex according to claim 12 of the formula:



wherein,

V is vitamin B₁₂ or a vitamin B₁₂ analogue, or derivative, bonded to X either through a carboxylate group pendant to the corrin nucleus of VB₁₂ or through the central cobalt
35 atom or to a functional group introduced onto the VB₁₂ molecule,

X is selected from: -NHNH-, -NH-, -O-, S-, -SS- or -CH₂-, and A is an optionally

substituted, saturated or unsaturated, branched or linear, C₁₋₅₀ alkylene, cycloalkylene or aromatic group, optionally with one or more carbons within the linear chain being replaced with N, O or S, and wherein the optional substituents are selected from carbonyl, carboxy, hydroxy, amino and other groups, and

Y is the covalent linkage between A and Z where Y is selected from -NHCO-, -CONH-, -CONHNHCO-, -N=N-, -N=CH-, -NHCH₂-, -NHN=CH-, -NHNHCH₂-, -SS-, -SCH₂-, -CH₂S-, -NHCRNH-, [R is O, S or NH₂], -COO-, -OCO-, and Z is EPO.

15. A complex according to claim 12 wherein said vitamin B₁₂ analogue is selected from cyano-cobalamin (CN-Cbl), aquocobalamin, adenosylcobalamin, methylcobalamin, hydroxy-cobalamin, 5-methoxybenzyl(cyano)cobalamin [(5-MeOBz)CN-Cbl]; the desdimethyl, monoethylamide and the methylamide analogues of these; alkyl cobalamins in which the alkyl chain is linked to the corrin nucleus by a direct Co C covalent bond; chlorocobalamin, sulfitecobalamin, nitrocobalamin, thiocyanatocobalamin, benzimidazole(cyano)cobalamin derivatives including 5,6-dichlorobenzimidazole, 5-hydroxybenzimidazole, trimethylbenzimidazole; adenosylcyanocobalamin [(Ade)CN-Cbl], cobalamin lactone, cobalamin lactam; the anilide, ethylamide, monocarboxylic and dicarboxylic acid derivatives of VB₁₂ or its analogues; the mono-, di- and tricarboxylic acid derivatives or the propionamide derivatives of eVB₁₂; and analogues of VB₁₂ in which the cobalt is replaced by zinc or nickel.

16. A complex according to claim 12 wherein said vitamin B₁₂ analogue is an analogue where the corrin ring is substituted with a substituent which does not affect binding to intrinsic factor.


17. A complex according to claim 12 wherein said EPO is covalently bound to a pharmaceutically acceptable polymer.

18. The complex according to claim 17 wherein the polymer selected from dextran, inulin, cellulose, starch and derivatives thereof, chondroitin sulfate, poly[N- α -(2-hydroxypropyl)-methacrylamide] and derivatives thereof, styrene-maleic anhydride copolymer, divinylether-maleic anhydride copolymer, polylysine, poly(glutamic acid), poly(hydroxypropyl glutamine), poly(lactic acid), water soluble polyurethanes formed by covalent linkage of PEG with lysine or other amino acids and branched chain polypeptides.
19. A complex according to claim 17 wherein said pharmaceutically acceptable polymer is biodegradable within the human or animal body.
20. A process for the production of a complex between EPO and a carrier selected from vitamin B₁₂ or a vitamin B₁₂ analogue which comprises: reacting one or both of said carrier and EPO with a diradical spacer having terminal reactive groups to form a carrier/linker and/or EPO linker intermediate, and thereafter reacting together the respective components to give a complex between EPO and said carrier wherein the components are covalently linked through a said spacer.
21. A process according to claim 20 wherein EPO and/or said carrier are modified prior to derivatization to provide at least one functional group capable of forming a chemical linkage.
22. A process according to claim 20 wherein said carrier is covalently linked to EPO through: an amide linkage, an acyl hydrazide, an imine linkage, or a hydrazone linkage.
23. A reagent for the detection of buried thiol groups of a protein or peptide, said reagent comprising a complex of either vitamin B₁₂ (or an analogue thereof) or more generally of a label detectable by visual or instrumental methods which is covalently linked to a diradical spacer, said spacer having a terminal reactive group capable of forming a

covalent bond with a free thiol in said protein or peptides.

24. A reagent according to claim 23, wherein said spacer has a backbone of from 1-50 atoms.

25. A reagent according to claim 23, of the general formula:



wherein,

V' is vitamin B₁₂ or a vitamin B₁₂ analogue, bonded to X either through a carboxylate group pendant to the corrin nucleus of VB₁₂ or through the central cobalt atom or to a functional group introduced onto the VB₁₂ molecule, or V' is a label detectable by visual or instrumental methods, and

X is selected from: -NHCO-, -CONH-, -CONHNHCO-, -N=N-, -N=CH-, -NHCH₂-, -NHN=CH-, -NHNHCH₂-, -SS-, -SCH₂-, -CH₂S-, -NHCRNH-, [R is O, S or NH₂], -COO-, -OCO-, and

A is an optionally substituted, saturated or unsaturated, branched or linear, C₁₋₅₀ alkylene, cycloalkylene or aromatic group, optionally with one or more carbons within the chain being replaced with N, O or S, and wherein the optional substituents are selected from carbonyl, carboxy, hydroxy, amino and other groups, and

Y' is a functional group capable of reacting with thiols to give a stable covalent linkage.

26. A reagent according to claim 25 wherein Y' is selected from iodoacetyl, bromoacetyl, chloroacetyl, maleimido, 3-carboxy-4-nitrophenyldithio or 2-pyridyldithio.

27. A reagent according to claim 23 wherein said label is selected from chromagens, catalysts, fluorescent compounds, chemiluminescent compounds, radioactive isotopes, colloidal metal and non-metallic particles, dye particles, enzymes or substrates, antibodies or antigens, biotin, avidin or streptavidin, latex particles, liposomes or other vesicles containing signal producing substances.

28. A pharmaceutical composition which comprises a complex according to claims 1 to 8 or 12 to 19 optionally in association with a pharmaceutically acceptable carrier or excipient.

5

29. A method for treating disease in a patient responsive to therapy with GCSF or EPO which comprises administering to said patient a therapeutically effective amount of a complex according to any one of claims 1 to 8 or 12 to 19, or a composition according to claim 28.

10

30. Use of a complex according to any one of claims 1 to 8 or 12 through 19 for the manufacture of a medicament

15 31. Use of a complex according to any one of claims 1 to 8 or 12 through 19 for the manufacture of a medicament.

AMENDED CLAIMS

[received by the International Bureau on 7 November 1994 (07.11.94);
original claims 23-27 amended; remaining claims unchanged (2 pages)]

18. The complex according to claim 17 wherein the polymer selected from dextran, inulin, cellulose, starch and derivatives thereof, chondroitin sulfate, poly[N- α -(2-hydroxypropyl)-methacrylamide] and derivatives thereof, styrene-maleic anhydride copolymer, divinylether-maleic anhydride copolymer, polylysine, poly(glutamic acid),
5 poly(hydroxypropyl glutamine), poly(lactic acid), water soluble polyurethanes formed by covalent linkage of PEG with lysine or other amino acids and branched chain polypeptides.
19. A complex according to claim 17 wherein said pharmaceutically acceptable polymer
10 is biodegradable within the human or animal body.
20. A process for the production of a complex between EPO and a carrier selected from vitamin B₁₂ or a vitamin B₁₂ analogue which comprises: reacting one or both of said carrier and EPO with a diaradical spacer having terminal reactive groups to form a
15 carrier/linker and/or EPO linker intermediate, and thereafter reacting together the respective components to give a complex between EPO and said carrier wherein the components are covalently linked through a said spacer.
21. A process according to claim 20 wherein EPO and/or said carrier are modified prior to
20 derivatization to provide at least one functional group capable of forming a chemical linkage.
22. A process according to claim 20 wherein said carrier is covalently linked to EPO through: an amide linkage, an acyl hydrazide, an imine linkage, or a hydrazone linkage.
25
23. A method for the detection of buried thiol groups of a protein or peptide, said method comprising the use of a reagent such as a complex of either vitamin B₁₂ (or an analogue thereof) or more generally of a label detectable by visual or instrumental methods which is covalently linked to a diradical spacer, said spacer having a terminal reactive group
30 capable of forming a

AMENDED SHEET (ARTICLE 19)

covalent bond with a free thiol in said protein or peptides.

24. The method according to claim 23, wherein said spacer has a backbone of from 1-50 atoms.

25. The method according to claim 23 wherein said reagent is of the general formula:



wherein,

V' is vitamin B₁₂ or a vitamin B₁₂ analogue, bonded to X either through a carboxylate group pendant to the corrin nucleus of VB₁₂ or through the central cobalt atom or to a functional group introduced onto the VB₁₂ molecule, or V' is a label detectable by visual or instrumental methods, and

X is selected from: -NHCO-, -CONH-, -CONHNHCO-, -N=N-, -N=CH-, -NHCH₂-, -NHN=CH-, -NHNHCH₂-, -SS-, -SCH₂-, -CH₂S-, -NHCRNH-, [R is O, S or NH₂], -COO-, -OCO-, and

A is an optionally substituted, saturated or unsaturated, branched or linear, C₁₋₅₀ alkylene, cycloalkylene or aromatic group, optionally with one or more carbons within the chain being replaced with N, O or S, and wherein the optional substituents are selected from carbonyl, carboxy, hydroxy, amino and other groups, and

Y' is a functional group capable of reacting with thiols to give a stable covalent linkage.

26. The method according to claim 25 wherein Y' is selected from iodoacetyl, bromoacetyl, chloroacetyl, maleimido, 3-carboxy-4-nitrophenyldithio or 2-pyridyldithio.


27. The method according to claim 23 wherein said label is selected from chromagens, catalysts, fluorescent compounds, chemiluminescent compounds, radioactive isotopes, colloidal metal and non-metallic particles, dye particles, enzymes or substrates, antibodies or antigens, biotin, avidin or streptavidin, latex particles, liposomes or other vesicles containing signal producing substances.

AMENDED SHEET (ARTICLE 19)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU 94/00274

A. CLASSIFICATION OF SUBJECT MATTER Int. Cl. ⁵ A61K 31/68, 37/02, 47/48 According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A61K 37/02: GRANULOCYTE COLONY STIMULATING FACTOR OR GCSF ERYTHROPOIETIN OR EPO Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base, and where practicable, search terms used) WPAT, CASM JAPIO				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.		
X	WO 87/02251 (BIOTECHNOLOGY AUSTRALIA PTY LTD) 23 April 1987 (23.04.87)	23		
X A	DE,A, 2546476 (ISRAEL, MURRAY) 21 April 1977 (21.04.77)	23 1-22, 24-43		
Y	WO 93/25221 (ALKERMES CONTROLLED THERAPEUTICS, INC) 23 December 1993 (23.12.93)	17, 18		
Y	US 5206219 (APPLIED ANALYTICAL INDUSTRIES, INC) 27 April 1993 (24.04.93)	17, 18		
<div style="display: flex; justify-content: space-between; align-items: flex-start;"> <div style="width: 45%;"> <input type="checkbox"/> Further documents are listed in the continuation of Box C. </div> <div style="width: 45%;"> <input checked="" type="checkbox"/> See patent family annex. </div> </div>				
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top;"> * Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed </td> <td style="width: 50%; vertical-align: top;"> "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family </td> </tr> </table>			* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family			
Date of the actual completion of the international search 23 August 1994 (23.08.94)	Date of mailing of the international search report 6 Sept 1994 (06.09.94)			
Name and mailing address of the ISA/AU AUSTRALIAN INDUSTRIAL PROPERTY ORGANISATION PO BOX 200 WODEN ACT 2606 AUSTRALIA Facsimile No. 06 2853929	Authorized officer  J. HANSON Telephone No. (06) 2832263			

International application No.
PCT/AU 94/00274

Patent Document Cited in Search Report		Patent Family Member	
WO	8702251	AU	65289/86
CA	1330791	EP	220030
DE	2546464	DE	2546474
WO	9325221	AU	46308/93
US	5206219	US	5206219
<p>END OF ANNEX</p>			

THIS PAGE BLANK (USPTO)